

### **Nanostictis**

On Lobaria. British Columbia. .... N. christiansenii Etayo

Etayo, J. and P. Diederich. 1996. Lichenicolous fungi from the western Pyrenees, France and Spain. III. Species on Lobaria pulmonaria. Bulletin de la societe des naturalistes luxembourgeois 97: 93-118.

**Nectria**

On Peltigera. British Columbia, Oregon, Alaska. .... N. lecanodes Cesati

## Nectriella

After Hawksworth, 1980, 1983, and Dennis

Perithecia subglobose, smooth, brightly colored, erumpent; otherwise like those of Nectria.

Ascospores (11-)12-15(-17) x (4-)4.5-7 um, hyaline, 1-septate, elliptic-cylindric, only slightly constricted at septum, smooth, containing several oil drops; to 8 per ascus, biseriate; Ascomata immersed within host tissue, then splitting it to expose their apices, ovoid, smooth, to 0.25 mm across, light red, brownish red to pale orange. Asci clavate, to 85 x 13 um. Anamorph Illosporium carneum. On Peltigera thalli. ....N. robergii (Mont. & Desm.)

Weese

**Neolamya**

On Peltigera. British Columbia. .... N. peltigerae (Mont.) Theiss. & Syd.

## Nesolechia

After Keissler, and Dennis

Apothecia sessile, erumpent, black, subgelatinous; asci 8-spored; ascospores hyaline, non-septate; paraphyses forked; hymenium I+ blue.

Spores almost spherical. Hypothecium brown, K-. On Parmelioid lichens. Florida (reported by Keissler, 1933). ..... N. thallicola

N. oxyspora = Phacopsis oxyspora

**Niesslia**

On Cladonia. British Columbia. .... N. cladoniicola D. Hawksw. & W. Gams

**Nigropuncta**

On crustose lichens. British Columbia. .... N. rugulosa D. Hawksw.