

Arthrorhaphis Th. Fr.
(ARTHRORHAPHIDACEAE)

After Poelt & Vezda, 1977, Purvis, 1992, and Ihlen 1998

Rev. 5/94

Thallus crustose, without a distinct prothallus, enclosed within the thallus of the host and gray to gray-green, or free at some stage and soon verrucose-squamulose, the squamules strongly convex, bright greenish yellow to whitish gray, ecorticate or with a colorless epinecral layer; soredia occasionally present. Photobiont chlorococcoid, forming a thick layer.

Apothecia sessile or located between the squamules, black, urceolate or discoid, mostly \pm filled with brownish green, crumbling, drop-like concretions. Discs flat, strongly marginate, but true exciple poorly developed, of lax hyphae with strongly swollen walls. Paraphyses free, thin, branched and net-like, anastomosing, the apices not or only weakly thickened. Hymenium containing oil droplets. Asci 8-spored, clavate to elongate-clavate or cylindrical, K/I-, the wall scarcely thickened at the apex, with an ocular chamber. Spores cylindrical to fusiform or mostly acicular, (2?-)7-12-septate.

Conidiomata unknown. On acidic substrates, either parasitic on Baeomyces species or, at least later, free-living on soil amongst mosses or on weathered rock surfaces in cool regions (arctic-alpine).

This genus is a segregate of Bacidia, distinguished especially by its bright yellowish-greenish thallus.

1. Spores 3(-5)-septate, 16-20 x 3-4 μ m, spindle-shaped; bordering verrucules of thallus effigurate, esorediate; hymenium and exciple green-tinged. Greenland. [see Thomson 1997 for full description]. A. vacillans

1. Spores over 6-septate, over 20 μ m long, needle-shaped; bordering verrucules of thallus sorediate or not; hymenium green-tinged or not. 2

2. Thallus of bullate-high convex composite squamules, not sorediate, up to 1 mm (or more) wide (but sometimes abraded). Medulla with calcium oxalate. Apothecia not always occurring, sometimes in rows, to 0.6(-1.5) mm or more wide. Spores (24-)29.5-41.7-54(-72) x (2.5-)4-5.4-7(8.5) μ m, Squamules \pm rounded, either remaining \pm discrete or becoming confluent and forming small, \pm compact irregular patches to ca. 1 cm diam.; marginal lobes indistinct, not becoming elongate; surface yellow to bright yellow-green, matt, often roughened and appearing \pm pruinose; heteromerous; cortex present; epinecral layer 7-10 μ m thick, continuous. Photobiont cells (7-)12-14(-18) μ m diam., \pm globose. Photobiont layer 60-70 μ m, continuous. Medulla with calcium oxalate crystals. Apothecia located between squamules, 0.1-0.6 mm diam. slightly concave, then flat, black; margin distinct, often raised, or becoming excluded; exciple brownish in the outer part, greenish in the inner part, (12-)25-55 μ m thick; hypothecium hyaline to green, K+ brown; true exciple and upper part of hymenium blackish gray-green. Hymenium hyaline to light green, with oil drops, K+ brown. Paraphyses branched, 1.5-2.5 μ m thick, not swollen at apices. Asci 80-110 x 12-17 μ m, cylrindrical to elongate-clavate, Spores arranged in parallel, narrowly fusiform or acicular, smooth, (6-)8-11(-14) septate.

Containing rhizocarpic acid and epanorin. On acid soil or decaying mosses, or on Baeomyces, often in crevices on \pm vertical siliceous rock faces, arctic-alpine.A. alpina

2. Thallus of flat to slightly convex 0.5 mm wide, soon sorediate erupting composite areoles, frequently very thin, fragile (readily broken when touched with a needle). Thallus lacking calcium oxalate. Apothecia to about 0.1 mm wide, often lacking. Spores (35-)47.5-61.0-74.5(-100) x (2-)3-4.4-5.5(-6) μ m. Without a distinct, continuous epinecral layer. Spores arranged in parallel, narrowly fusiform or acicular, smooth, (6-)8-13(-15) septate, Thallus yellow to yellowish green or bright greenish; areoles convex, partly or completely sorediate. Calcium oxalate crystals absent. Photobiont cells globose, 9-12 μ m. Apothecia black, 0.2-0.6 mm diam., disc flat to concave; exciple distinct, raised, greenish brown in outer part, green in inner part, 50-70 μ m thick. Hypothecium hyaline, K+ light brown. Hymenium green, with oil drops, K+ brown. Paraphyses branched, 1.5-2.5 μ m, not swollen at apices. Asci cylindrical to elongate-clavate, K/I-, (60-)75-110(-120) x 7-13 μ m. With rhizocarpic acid and epanorin. Similar habitats to A. alpina, often parasitic on Baeomyces or Dibaeis on acid soils and weathered lime-free rock.A. citrinella

Literature

Galloway, D. 1985. Flora of New Zealand Lichens.

Ihlen, P. G. 1998. The lichenicolous fungi on species of the genera Baeomyces, Dibaeis and Icmadophila in Norway. Lichenologist 30: 27-57.

Poelt, J. and A. Vezda. 1977. Bestimmungshclüssel europäischen Flechten. Erg. I. Cramer, Vaduz.

Purvis, W. 1992. Arthrorhaphis. In: Purvis, et al., Lichen Flora of Great Britain and Ireland.